

Front panel

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS

Racumin[®]

Domestic Rat and Mouse Paste 

Active Constituent: **0.37 g/kg COUMATETRALYL**

(rat/mice pictograms)

READY TO USE

200 g NET (20 x 10g bait sachets)

Back panel

ALL RODENTICIDES CAN BE HARMFUL TO PETS. DO NOT ALLOW THEM TO GAIN ACCESS TO THIS PRODUCT.

DIRECTIONS FOR USE

Restraints -

Do NOT place baits in open unless in bait stations.

Do NOT place baits in any position accessible to children, livestock, or domestic pets.

Do NOT remove bait from the paper sachet

SITUATION	PEST	RATE	CRITICAL COMMENTS
Domestic, (houses, homesteads, garden sheds, outbuildings) Urban and semi-urban commercial premises	Mice and Rats	3-6 sachets per bait site	Place bait sachets in areas of known rodent activity. For rats place a minimum of 3 sachets (30g) per placement. Do not exceed 3 m between baits for mice, or 9 m for rats. Follow the Baiting Strategy detailed below.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.

GENERAL INSTRUCTIONS: Do not apply directly on or near exposed food or food utensils, food preparation areas, livestock feed or water, food storage areas, or onto packaging materials or in areas where such packaging is stored. Do not feed the bait to wild or domestic birds.

Baiting Strategy

Eliminate, as far as practicable, all alternative food sources. Place the bait sachets in areas of known rodent activity and under cover (eg, behind appliances and furniture, in ceiling voids etc). Inspect baits daily. If eaten quickly increase the number of bait sachets. Continue observation and replenishment until no more bait is taken. Baiting for at least 2 weeks is necessary to reduce mouse or rat numbers. Caution is required in use of this product in and near piggeries. Do NOT allow to contaminate pig food.

PRECAUTION: Should not be used in areas accessible to children.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:

Place baits in locations inaccessible to other animals and birds. This product is hazardous to pigs, cats, dogs and poultry. Do not contaminate ponds, waterways or drains with the product, bait refuse from bait stations, or used containers.

STORAGE AND DISPOSAL:

Store in closed, original container in a safe place. Destroy untaken baits. Dispose of used containers by wrapping with paper and putting in garbage. Any spillage of bait should be removed immediately.

SAFETY DIRECTIONS:

Poisonous if swallowed. Do not touch the bait. If on skin, and after each baiting, wash thoroughly with soap and water. Vitamin K1 (Phytomenadione) is antidotal.

Back panel continued

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131126. If swallowed and if more than 15 minutes from a hospital, induce vomiting preferably using Ipecac Syrup APF.

Notice

Purchaser assumes all risk of use and handling where the product is not used in accordance with the directions given.

NRA Approval No. 52343/0400


BAYER AUSTRALIA LTD Racumin® is a trademark of Bayer
ACN 000 138 714
875 Pacific Highway, PYMBLE NSW 2073.



[B]

[DOM]

(label code)



Teabag

Issue Date: 18-6-01

Page: 4 of 4

The information above this line is not included in the label

Racumin[®] Paste

Coumatetralyl 0.375 g/kg

