

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Responsar[®] Beta SC

INSECTICIDE

Active Constituent: 125 g/L BETACYFLUTHRIN

GROUP 3A INSECTICIDE

For the control of cockroaches, spiders, ants, fleas, flies, silverfish, bedbugs, mosquitoes, clothes moths, carpet beetles and stored product pests in the situations specified in the DIRECTIONS FOR USE table.

GENERAL INSTRUCTIONS

Do not spray into air, directly on humans, exposed food or food utensils, food preparation areas, livestock feed or water, food storage areas or onto packing materials or in areas where such packaging is stored. When used in animal houses do not spray directly onto livestock, their food, litter, feed bins, or surfaces that are accessible to the stock. Use in food processing areas is limited to crack, crevice and barrier strip treatment only. Do NOT mist or fog this product.

Insecticide Resistance Warning

For insecticide resistance management Responsar is a Group **3A** insecticide. Some naturally occurring insect biotypes resistant to Responsar and other Group **3A** insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Responsar or other Group **3A** insecticides are used repeatedly. The effectiveness of Responsar on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from failure of Responsar to control resistant insects. Responsar may be subject to specific resistance management strategies. For further information contact your local supplier, Bayer CropScience representative.

Mixing

Add the required quantity of Responsar SC Insecticide to water while stirring.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

Dangerous to fish. Cover fish tanks before using. Do not contaminate ponds, waterways or drains with this product or used container.

STORAGE AND DISPOSAL, PROTECTION OF OTHERS

Store in tightly closed, original container in a safe, well ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal, and add rinsings to the tank mix or dispose of in a disposal pit away from waterways. Destroy empty containers by breaking, crushing or puncturing them. Bury containers at a depth of 500 mm or more at a safe disposal site, or take them to a dump that does not burn its refuse. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact, inhaled, or swallowed. Avoid contact with eyes and skin. Do not inhale vapour and spray mist. When preparing the spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and half face-piece respirator. When using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat, and rubber gloves. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, respirator and if rubber wash with detergent and warm water, and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone 131126. Remove from contaminated area. Apply artificial respiration if not breathing.

Additional Information is listed in the Material Safety Data Sheet.

LIMITATION OF LIABILITY

This product is designed only for the particular purposes indicated on this label. Failure to use this product strictly as directed may be illegal; prove dangerous; render the product ineffective and avoid any Warranty or Liability of Bayer CropScience Pty Ltd.

Resposar® is a Registered Trademark of Bayer

NRA Approval No 52198/0999

FOR 24 HOUR SPECIALIST ADVICE
IN EMERGENCY ONLY
PHONE 1800 033 111

DIRECTIONS FOR USE (all states):

SITUATIONS	RATE	PEST	CRITICAL COMMENTS
Meatworks (non-product areas), Farm buildings, Food processing establishments, Shops, Offices, Factories, Storerooms, Hospitals, Schools, Barracks, Houses, Ships, Rubbish Tips	Initial or Cleanout rate 4 mL/L Maintenance rate or when used as a tank mix with a larvicide (e.g. Starycide) 2 mL/L	Cockroaches	Thoroughly spray hiding places (eg. cracks and crevices, behind stoves and sinks, in cupboards, around drains, rubbish bins etc.) until point of run-off. If hiding places are not directly accessible, spray barrier strips around them. Repeat treatment as required.
		Spiders	Thoroughly spray spiders, their webs and crevices in which they hide. Do not remove webs for a few days.
		Ants	Where possible locate the nest and spray. If nest is inaccessible, spray all infested areas.
		Fleas	Clean or vacuum infested areas and burn all dust collected. Lightly spray all infested areas especially under furniture. Pets should be treated with a suitable flea control product registered for use on animals.
		Flies Silverfish	Spray surfaces where flies are known to rest eg. on ceilings or around windows. Effective for many weeks.
			Locate the source of the infestation and spray. If it cannot be located spray all surfaces over which silverfish travel, especially inside cupboards.
		Clothes moths	Spray clothes storage areas.
		Carpet beetles	Lightly spray carpets, particularly in areas of little or no traffic.
		Bedbugs	Fumigate all infested mattresses or clothing. Wash sheets and clothing before re-use. Apply Resposar SC to walls, cracks and crevices, skirting boards and bed frames.
		Mosquitoes	Apply as a wet spray till point of run-off, to areas where mosquitoes rest such as walls and awnings, fences and some vegetated areas.
Stored Product Pests	Clean and vacuum infested areas and burn all material collected. Thoroughly spray infested areas taking care to treat all cracks and crevices where pests may hide.		
Rubbish tips	Knockdown spray 1 mL/2 L	Flies	Spray surfaces of rubbish. Apply 200-300 mL of spray/m ² .

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.
USE IN FOOD PROCESSING AREAS IS LIMITED TO CRACK, CREVICE AND BARRIER STRIP TREATMENT ONLY.
TO BE USED BY LICENSED PEST CONTROL MANAGERS ONLY**