

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Sumilarv[®]

INSECT GROWTH REGULATOR

Active Constituent: 20 g/L PYRIPROXYFEN

GROUP 7C INSECTICIDE

For use with an adulticide to control cockroaches and fleas in domestic, industrial and public health situations as per the DIRECTIONS FOR USE table. Breaks the breeding cycle.

TO BE USED BY LICENSED COMMERCIAL PEST CONTROL OPERATORS ONLY

GENERAL INSTRUCTIONS

Insecticide Resistance Warning

For insecticide resistance management, Sumilarv Insect Growth Regulator is a Group 7C insecticide. Some naturally occurring insect biotypes resistant to Sumilarv and other Group 7C insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Sumilarv or other Group 7C insecticides are used repeatedly. The effectiveness of Sumilarv on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from the failure of Sumilarv to control resistant insects. For information about strategies for managing insecticide resistance contact your local supplier or Bayer Environmental Science representative.

Where to Apply

Cockroaches – areas where cockroaches harbour, paying particular attention to cracks and crevices.

Fleas - skirting boards, floors and carpets and adjacent wall area.

Do not spray the moving parts of any machinery, electric motors or switchgear with the water-based product.

PRECAUTIONS

AVOID CONTACT WITH FOOD, FOOD UTENSILS, OR PLACES WHERE FOOD IS PREPARED OR STORED.

BEFORE USE - REMOVE OR COVER ALL EXPOSED FOODSTUFFS. COVER ALL DISHES AND UTENSILS, AND PLACES WHERE FOOD IS PREPARED OR STORED.

AFTER USE - THOROUGHLY VENTILATE TREATED AREA. CLEAN UP THOROUGHLY BEFORE PROCESSING/SERVING RESUMES.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and other aquatic life. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray wear elbow length PVC gloves. Wash hands after use. After each day's use wash gloves.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (telephone 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

Sumilarv[®] is a Registered Trademark of Sumitomo Chemical Co. Ltd.
Cislin[®] and Crackdown[®] are Registered Trademarks of Bayer

APVMA Approval No.: 500/1103

FOR 24 HOUR SPECIALIST ADVICE
IN EMERGENCY ONLY
PHONE 1800 033 111

DIRECTIONS FOR USE

SITUATION	PEST	RATE	CRITICAL COMMENTS
Houses, flats, factories, shops, warehouses, restaurants, public buildings	Fleas	<u>Initial treatment</u> 25 mL / 5 L water applied as a tank mix with the label rate of a registered adulticide such as Cislin[®] or Crackdown[®]	Apply as a coarse spray at the rate of 5 L / 100 m ² or to the point of run off which will vary with the nature of the surface being treated. Re-apply when signs of infestation reappear.
	Cockroaches	<u>Maintenance treatments</u> 25 mL / 5 L water applied alone 50 mL / 5 L water applied as a tank mix with the label rate of a registered adulticide such as Cislin or Crackdown	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.